

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2008		2. REPORT TYPE		3. DATES COVERED	
4. TITLE AND SUBTITLE Lateral Mixing				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) APL/UW,1013 NE 40th Str,Seattle,WA,98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES The original document contains color images.					
14. ABSTRACT The major research goal for 2008 was to publish an analysis of the relative importance of diapycnal and isopycnal mixing on the Oregon Shelf and study the role of internal waves in this environment					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Lateral Mixing

Eric A. D'Asaro

APL/UW 1013 NE 40th Str, Seattle, WA 98105

phone: (206) 685-2982 fax: (206) 543-6785 email: dasaro@apl.washington.edu

Award Number: N00014-08-1-0974

<http://opd.apl.washington.edu/~dasaro/HOME/>

LONG-TERM GOALS

I seek to understand the processes controlling lateral mixing in the ocean, particularly at the submesoscale, i.e. 100m-20km.

OBJECTIVES

The major research goal for 2008 was to publish an analysis of the relative importance of diapycnal and isopycnal mixing on the Oregon Shelf and study the role of internal waves in this environment.

The major administrative goal for 2008 was to work with others to develop a plan for the Scalable Lateral Mixing DRI.

APPROACH

Neutrally buoyant Lagrangian floats were deployed on the Oregon Shelf during the summers of 2000 and 2001. The float motion measures the rate of change of water properties along parcel trajectories. High frequency measurements along the float trajectory were used to infer diapycnal mixing rates. The comparison of the observed rate of change and that inferred from diapycnal processes alone allows the role of isopycnal processes to be inferred.

Specifically, an equation for the rate of change of 'spice', Π , a passive scalar roughly orthogonal to potential density in θ/S space, can be derived roughly following McDougall (1984): $\Pi_t = D\Pi_{\sigma\sigma}\sigma_d^2$, where D is the diapycnal diffusivity, σ is potential density, and the subscripts σ and d imply differentiation with respect to potential density and depth respectively. The diapycnal diffusivity is evaluated from the spectrum of potential density near N following D'Asaro and Lien (2007) and the other quantities from the float CTD measurements. This analysis was applied to a single float from the 2001 Oregon Shelf float deployments.

A DRI planning meeting was held in May 2008 at MIT.

WORK COMPLETED

The results of the mixing analysis have been published in the Limnology and Oceanography special issue on Autonomous and Lagrangian Science.

A draft plan for the DRI was developed including two potential measurement programs and a mechanism for deciding on experimental sites.

RESULTS

Diapycnal mixing is found to mostly predict the observed changes at the float, with some notable periods where isopycnal mixing must also be important. Thus both can be important at the measured scales of many meters and a few days. A similar analysis conducted for chlorophyll instead of spice, shows that diapycnal mixing does not explain the observed variations. Instead, isopycnal mixing, planktonic sinking or possibly growth is important.

REFERENCES

D'Asaro, Eric, Ren-Chieh Lien, 2007, Measurement of Scalar variance dissipation from Lagrangian floats, , *J. Atmos. & Oceanic Tech.*, 24, 1066-1077

McDougall, T.J. 1984. The relative roles of diapycnal and isopycnal mixing on subsurface water mass conversion. *J. Phys. Oceanogr.* **14**: 1577–1589.

PUBLICATIONS

D'Asaro, Eric A. 2008. A diapycnal mixing budget on the Oregon shelf. *Limnol. Oceanogr.* 53: 2137-2150